

Supplementary Material

***In Silico* Transcriptomic Analysis of the Chloride Intracellular Channels (CLIC) Interactome Identifies a Molecular Panel of Seven Prognostic Markers in Patients with Pancreatic Ductal Adenocarcinoma**

Dimitrios E. Magouliotis¹, Nikos Sakellaridis², Konstantinos Dimas³, Vasiliki S. Tasiopoulou⁴, Konstantina A. Svokos⁵, Alexis A. Svokos⁶ and Dimitris Zacharoulis^{7,*}

¹Division of Surgery and Interventional Science, Faculty of Medical Sciences, UCL, London, UK and Department of Surgery, University of Thessaly, Biopolis, Larissa, Greece; ²Department of Pharmacology, Faculty of Medicine, School of Health Sciences, University of Thessaly, Biopolis, Larissa, Greece; ³Department of Pharmacology, Faculty of Medicine, School of Health Sciences, University of Thessaly, Biopolis, Larissa, Greece; ⁴Faculty of Medicine, School of Health Sciences, University of Thessaly, Biopolis, Larissa, Greece; ⁵The Warren Alpert Medical School of Brown University, Providence, RI, USA; ⁶Geisinger Medical Center, Danville, PA, USA; ⁷Department of Surgery, University Hospital of Larissa, Larissa, Greece

Figure S1. Flow chart




Figure S2. The interactome of CLIC-related genes, as produced in GeneMania Platform.






Figure S3. A. The median survival period was greater for PTGES2 high expression group, as compared with PTGES2 low expression group (HR: 0.6 [95% CI: 0.42, 0.87]; $p=0.007$). **B.** GSTP1 low expression group had an increased median survival period, as compared with the high expression group (HR: 1.41 [95% CI: 1.12, 1.78]; $p=0.004$).



A.



B.

Table S1. Clinical and pathological characteristics of the samples and patients include in the present study

STUDY ID, YEAR	GSE15471	GSE16515	GSE32676
Mean age at enrollement (years)	N/A	N/A	67
Females (%)	N/A	N/A	52
R0 resection (%)	N/A	N/A	84
I	N/A	N/A	0
IIA	N/A	N/A	0
IIB	N/A	N/A	0
III	N/A	N/A	8
IV	N/A	N/A	92